arXiv:1301.3567v2 [math-ph] 6 Feb 2013

Ermakov-Pinney Equations with Abel-induced Dissipation

Stefan C. Mancas*
Department of Mathematics,
Embry-Riddle Aeronautical University,
Daytona Beach, FL. 32114-3900, U.S.A.

Haret C. Rosu†
IPICyT, Instituto Potosino de Investigacion Cientifica y Tecnologica,
Apdo. Postal 3-74 Tangamanga, 78231 San Luis Potosí, S.L.P., Mexico

We introduce a dissipative Ermakov-Pinney system with the dissipation based on the corresponding integrable Abel equation. Based on a particular solution, the general solution to dissipative Ermakov-Pinney equations of this type is constructed, including the cases of higher-order Reid nonlinearities, by using the Ermakov invariant of the system.

I. INTRODUCTION

The nonlinear Ermakov-Pinney (EP) equation has long been known to have profound connections with the linear equations of identical operatorial form without the inverse cubic nonlinearity and because of this it has been seen as an example of 'nonlinearity from linearity' [1]. A historical overview has been written by Leach and Andriopoulos [2] and the fundamental importance of the EP equation for parametric oscillators, both classical and quantum-mechanical, with their vast application reaches, is well established in the literature.

As known, in 1880, Ermakov introduced the following system [3]

$$\begin{cases} \ddot{y}+\omega^2(t)y=0\ , \\ \ddot{x}+\omega^2(t)x-\kappa x^{-3}=0\ , \qquad \kappa>0\ , \end{cases} \tag{1}$$

which is endowed with the so-called Ermakov dynamic invariant

$$I_{0-\kappa}(x,y,\dot{x},\dot{y})=\kappa\left(\frac{y}{x}\right)^2+(x\dot{y}-y\dot{x})^2\ . \tag{2}$$

We label here this invariant by the tuning parameters of the nonlinear term of the two equations.
A century later, Ray and Reid [4] and also Sarlet [5] introduced generalized Ermakov systems and their associated invariants of the type $I_{f,g}$, where f and g are functions of the quotient variable y/x. Recent works on the connections of the Ermakov systems with the nonlinear superposition principle belong to Cariñena and collaborators [6]. When $f=g=-\kappa$ and one knows a particular solution x_p, the invariant can be written in the following form

$$I_{-\kappa-\kappa}=\kappa\left[\left(\frac{x}{x_p}\right)^2+\left(\frac{x_p}{x}\right)^2\right]+(x_p\dot{x}-\dot{x}_px)^2\ . \tag{3}$$

In this paper, we start by shortly presenting the factorization of linear dissipative ODE which we use to introduce a special EP equation with dissipation determined by an integrability condition for the Abel equation of the first kind [7]. Next, we include this special Ermakov-Pinney equation in a new type of Ermakov system and determine its general solution by means of the attached invariant.

II. LINEAR DISSIPATIVE ODES AND THEIR FACTORIZATION

Linear dissipative ODEs of the type

$$u_{\zeta\zeta}+g(\zeta)u_\zeta+h(\zeta)u=0 \tag{4}$$

---

[*] Electronic address: stefan.mancas@erau.edu
[†] Electronic address: hcr@ipicyt.edu.mx

can be factored similarly to their nonlinear counterparts [7], i.e.,

$$\left(\frac{d}{d\zeta}-\Phi_2\right)\left(\frac{d}{d\zeta}-\Phi_1\right)u=0\ , \tag{5}$$

which leads to

$$u_{\zeta\zeta}-\frac{d}{d\zeta}\left(\Phi_1(\zeta)u\right)-\Phi_2u_\zeta+\Phi_1\Phi_2u=0\ . \tag{6}$$

Furthermore, one gets

$$u_{\zeta\zeta}-\left(\Phi_1+\Phi_2\right)u_\zeta+\left(\Phi_1\Phi_2-\Phi_1'\right)u=0\ , \tag{7}$$

where the prime stands for $\frac{d}{d\zeta}$.

Then, we have

$$\begin{cases} g(\zeta)= & -(\Phi_1+\Phi_2)\\ h(\zeta)= & \Phi_1\Phi_2-\Phi_1'\ , \end{cases}$$

which is a system of equations slightly different from that first used by Rosu and Cornejo-Pérez [8] for nonlinear equations of the same structure as (4). This system leads immediately to the following Riccati equation

$$\Phi_1'=-\Phi_1^2-g\Phi_1-h\ . \tag{8}$$

Moreover, since we can always reduce the dissipative ODE (4) to a canonical form without dissipation via transformation $u(\zeta)=\hat{u}(\zeta)e^{-\frac{1}{2}\int g(\zeta)d\zeta}$, then we will obtain

$$\hat{u}_{\zeta\zeta}+\hat{h}(\zeta)\hat{u}=0, \tag{9}$$

where $\hat{h}(\zeta)=h(\zeta)-\frac{1}{4}g^2(\zeta)-\frac{1}{2}\frac{d}{d\zeta}g(\zeta)$, then, without loss of generality, we will always solve the Riccati equation (8) with $g(\zeta)=0$.

From (5), we also have

$$u_\zeta=\Phi_1u \tag{10}$$

which leads to

$$\frac{du}{u}=\Phi_1d\zeta \tag{11}$$

from where

$$u=c_1\,e^{\int^\zeta\Phi_1(x)dx}\ . \tag{12}$$

In other words, for all functions $h(\zeta)$ for which the Riccati equation is integrable, one first determines the factoring function Φ_1, and writes the particular solution of (4) as the exponential in (12).

III. EP EQUATION AS A LINEAR DISSIPATIVE ODE

Take now the EP equation:

$$u_{\zeta\zeta}+p(\zeta)u+u^{-3}=0, \tag{13}$$

and consider it as a linear dissipative ODE of the type (4), so $g(\zeta)=0$ and $h(\zeta)=p(\zeta)+cu^{-4}$. The corresponding Riccati equation is

$$\Phi_1'=-\Phi_1^2-h \tag{14}$$

and from (12), one can write

$$h(\zeta) = p(\zeta) + c\left(c_1 e^{\int^{\zeta}\Phi_1(x)dx}\right)^{-4} \ . \tag{15}$$

As is well known, if we have two linear independent solutions u_1, and u_2 to

$$u_{\zeta\zeta} + h(\zeta)u = 0, \tag{16}$$

then the solution of the corresponding EP equation

$$v_{\zeta\zeta} + h(\zeta)v + cv^{-3} = 0 \tag{17}$$

is given by [9]

$$v(\zeta) = \sqrt{u_1^2 - \frac{cu_2^2}{W^2}} \ , \tag{18}$$

where W is the Wronskian of the two linearly independent solutions.

A particular solution to (16) that will be later used in finding a general solution can be obtained as follows: Let's take $h(\zeta) = \lambda^2 > 0$, a constant, which is a case corresponding to the simple harmonic oscillator equation

$$u_{\zeta\zeta} + \lambda^2 u = 0, \tag{19}$$

with solutions $u_1 = \cos\lambda\zeta$ and $u_2 = \sin\lambda\zeta$ and Wronskian $W = \lambda$.

The EP equation for this case reads

$$v_{\zeta\zeta} + \lambda^2 v + cv^{-3} = 0 \tag{20}$$

and its solution can be written in the form

$$v = \sqrt{1 - \left(1 + \frac{c}{\lambda^2}\right)\sin^2\lambda\zeta} \ . \tag{21}$$

If we now use the factorization method from [7], with $g(v) = 0$ and $h(v) = \lambda^2 v + cv^{-3}$, the corresponding Abel equation for the EP equation (20) is:

$$\eta\frac{d\eta}{dv} + h(v) = 0 \ , \tag{22}$$

which gives

$$\eta^2 = c_1 - \lambda^2 v^2 + cv^{-2} \ . \tag{23}$$

We will see later that the integration constant c_1 is actually an invariant of (20).

IV. EP EQUATION WITH ABEL-INDUCED DISSIPATION

We build now the Ermakov-Pinney equation with Abel-type dissipation according to the equation

$$v_{\zeta\zeta} + g_v(v)v_\zeta + h_v(v) = 0 \tag{24}$$

proposed by us in a previous work [7], where g_v is obtained from h_v using Chiellini's integrability condition for Abel's equation, $\frac{d}{du}\left(\frac{h(u)}{g(u)}\right) = kg(u)$, k a real constant, which leads to

$$g_v = \frac{h_v}{\sqrt{c_1 + k(\lambda^2 v^2 - \frac{c}{v^2})}} \ . \tag{25}$$

Thus, one gets

$$g_v = \frac{\lambda^2 v^2 + cv^{-2}}{\sqrt{k\lambda^2 v^4 + c_1 v^2 - kc}} \ . \tag{26}$$

The inverse of the Abel solution, $\frac{1}{\eta}=\frac{g_v}{h_v}|_{k=-2}$, see [7], is obtained from

$$\frac{1}{\eta}=\frac{v}{\sqrt{-2\lambda^2v^4+c_1v^2+2c}}~, \tag{27}$$

which gives

$$\zeta-\zeta_0=\int\frac{vdv}{\sqrt{-2\lambda^2v^4+c_1v^2+2c}}~. \tag{28}$$

Let $\omega=\sqrt{2}\lambda v^2$, then (28) becomes

$$\zeta-\zeta_0=\frac{1}{2\sqrt{2}\lambda}\int\frac{d\omega}{\sqrt{a^2-\left(\omega-\frac{c_1}{2\sqrt{2}\lambda}\right)^2}}~, \tag{29}$$

where $a^2=c+\frac{c_1^2}{8\lambda^2}$, which gives the *particular* solution $v(\zeta)$, via

$$v^2=\frac{c_1\pm\sqrt{8\lambda^2c+c_1^2}\sin\left(2\sqrt{2}\lambda(\zeta-\zeta_0)\right)}{4\lambda^2}. \tag{30}$$

This equation is extremely important since the constant c_1 is an invariant [10].

V. GENERAL SOLUTION TO THE EP EQUATION WITH ABEL-INDUCED DISSIPATION

We will introduce now the following Ermakov system with Abel-induced dissipation

$$\begin{cases} u_{\zeta\zeta}+g_u(u)u(\zeta)+h(\zeta)u+bu^{-3}=0\\ v_{\zeta\zeta}+g_v(v)v(\zeta)+\lambda^2v+dv^{-3}=0~. \end{cases} \tag{31}$$

We have the following result.

Theorem. For the dissipative Ermakov system (31) with general $h(\zeta)$, and v a particular solution via (30) and g_v given by (26), the general solution $u(\zeta)$ is obtained by

$$u(\zeta)=\left[\frac{I_{bd}\pm\sqrt{I_{bd}^2+2bp}\sin\left(2\sqrt{p}(\Theta-\Theta_0)\right)}{2p}\,\frac{c_1\pm\sqrt{8\lambda^2c+c_1^2}\sin\left(2\sqrt{2}\lambda(\zeta-\zeta_0)\right)}{4\lambda^2}\right]^{\frac{1}{2}}~, \tag{32}$$

where I_{bd} is the following invariant

$$I_{bd}=-b\left(\frac{v}{u}\right)^2-d\left(\frac{u}{v}\right)^2+(u_\zeta v-uv_\zeta)^2. \tag{33}$$

Proof:

First, we construct the invariant I_{bd}, using (33).
Then, we let $\Theta=\int\frac{1}{v^2}d\zeta$, and $\theta=\frac{u}{v}$ in (33) to obtain the following separable equation

$$\frac{\theta d\theta}{\sqrt{b+I_{bd}\theta^2+d\theta^4}}=d\Theta~. \tag{34}$$

Since $d=-p<0$, then (34) has the same format as (28) with $p=2\lambda^2$, $I_{bd}=c_1$, $b=2c$. Then, by integrating we obtain

$$\theta^2=\frac{I_{bd}\pm\sqrt{I_{bd}^2+2bp}\sin\left(2\sqrt{p}(\Theta-\Theta_0)\right)}{2p}. \tag{35}$$

But from $\theta=\frac{u}{v}$, one gets $u=\theta v$ so that using (35) and (30) leads to (32).

We give now two related lemmas.

Lemma 1: The invariant I_{bd} is related to the factoring functions of the system (31).

Let Φ_1, and Ψ_1 be the factoring functions such that $u_\zeta = \Phi_1 u \to u = c_u e^{\int \Phi_1 d\zeta}$, and $v_\zeta = \Psi_1 v \to v = c_v e^{\int \Psi_1 d\zeta}$. Then, using (33) we have

$$I_{bd} = \theta^2 v^4 (\Phi_1 - \Psi_1)^2 - m\cosh\Big(\int (\Phi_1 - \Psi_1) d\zeta\Big) \ , \tag{36}$$

where $m = 2\Big(b\frac{c_v^2}{c_u^2} + d\frac{c_u^2}{c_v^2}\Big)$.

Lemma 2: $I = c_1$.

If one writes (33) as

$$I_{bd} = -b\frac{1}{\theta^2} - d\theta^2 + \Big(\frac{d\theta}{d\Theta}\Big)^2, \tag{37}$$

using chain rule, (35), and by substituting $\Theta = \int \frac{1}{v^2} d\zeta$ leads to $I_{bd} \equiv c_1$.
Let us choose $\lambda = c = \frac{1}{2}$, and $\zeta_0 = 0$ in (30), then

$$\Theta = \mp\sqrt{2}\mathrm{arctanh}\Big[\sqrt{1 + \mathrm{c}_1^2} \pm \mathrm{c}_1 \tan(\zeta/\sqrt{2})\Big], \tag{38}$$

and

$$\theta^2 = c_1 \pm \sqrt{c_1^2 + 1}\sin(\sqrt{2}\Theta). \tag{39}$$

By substituting both into (37), with $b = 1$, $d = -\frac{1}{2}$, and $\Theta_0 = 0$, then $I = c_1$.

VI. ABEL-DISSIPATIVE EP EQUATION WITH A $2m - 1$ NEGATIVE POWER NONLINEARITY

Reid has shown in [11] that the general solution to

$$v_{\zeta\zeta} + h(\zeta)v = q_m(\zeta)v^{-(2m-1)} \tag{40}$$

is

$$v = \big(u_1^m + \frac{\tilde{c}}{(m-1)W^2}u_2^m\big)^{1/m} \tag{41}$$

provided that u_1 and u_2 are two independent solutions of (16), and

$$q_m(\zeta) = \tilde{c}(u_1 u_2)^{m-2} \ . \tag{42}$$

We are interested in a general solution to

$$u_{\zeta\zeta} + g_u(u)u_\zeta + h(\zeta)u = q_m(\zeta)v^{1-2m}, \tag{43}$$

via the machinery of invariants.

First, let us choose $h(\zeta) = +\lambda^2, 0, -\lambda^2$, then (40) has solutions

$$\begin{cases} v_- = \big(a^m e^{m\lambda\zeta} + \tilde{c}_m b^m e^{-m\lambda\zeta}\big)^{1/m} \ , & h(\zeta) = -\lambda^2 \\ v_0 = (1 + \frac{\tilde{c}}{m-1}\zeta^m)^{\frac{1}{m}} \ , & h(\zeta) = 0 \ , \\ v_+ = \big(a^m \cos m\lambda\zeta + \tilde{c}_m b^m \sin m\lambda\zeta\big)^{1/m} \ , & h(\zeta) = \lambda^2, \end{cases} \tag{44}$$

where

$$\tilde{c}_m = \frac{\tilde{c}}{4\lambda^2 (ab)^m (m-1)} \ . \tag{45}$$

For simplification, let us write (44) as

$$\begin{cases} v_- = \left(Ae^{m\lambda\zeta} + Be^{-m\lambda\zeta}\right)^{1/m} \\ v_0 = (1 + B_0\zeta^m)^{\frac{1}{m}} \\ v_+ = \left(A\cos m\lambda\zeta + B\sin m\lambda\zeta\right)^{1/m} \ , \end{cases} \tag{46}$$

where $A = a^m$, $B = \frac{\tilde{c}}{4\lambda^2 a^m (m-1)}$, $B_0 = \frac{\tilde{c}}{m-1}$ and by integrating $\Theta_\mp = \int \frac{1}{v_\mp{}^2} d\zeta$ and $\Theta_0 = \int \frac{1}{v_0{}^2} d\zeta$ leads to

$$\begin{cases} \Theta_- = \frac{Ae^{2m\lambda\zeta}+B}{2\lambda B\left(Ae^{m\lambda\zeta}+Be^{-m\lambda\zeta}\right)^{2/m}} {}_2F_1\left(1, \frac{m-1}{m}; \frac{m+1}{m}; -\left(\sqrt{\frac{A}{B}}e^{m\lambda\zeta}\right)^2\right) \\ \Theta_0 = \zeta\, {}_2F_1\left(\frac{1}{m}, \frac{2}{m}; \frac{m+1}{m}; -B_0\zeta^m\right) \\ \Theta_+ = -\frac{\sin^{2/m}\left(m\lambda\zeta+\arctan{(A/B)}\right)\cos\left(m\lambda\zeta+\arctan{(A/B)}\right)}{m\lambda\left(A\cos m\lambda\zeta+B\sin m\lambda\zeta\right)^{2/m}} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} + \frac{1}{m}; \frac{3}{2}; \cos^2\left(m\lambda\zeta + \arctan{(A/B)}\right)\right) \ . \end{cases} \tag{47}$$

Finally, the solution to (43) is $u = v\theta$, with θ obtained from (39).
Using the same constants as above, together with $a = b = \tilde{c} = c_1 = 1$, then the solution is the following

$$\begin{cases} u_-(\zeta, m) = \left(e^{\frac{m\zeta}{2}} + \frac{1}{m-1}e^{-\frac{m\zeta}{2}}\right)^{\frac{1}{m}}\left(1 \pm \sqrt{2}\sin(\sqrt{2}\Theta_-)\right)^{1/2} \ , & h(\zeta) = -\lambda^2 \ , \\ u_0(\zeta, m) = \left(1 + \frac{1}{m-1}\zeta^m\right)^{\frac{1}{m}}\left(1 \pm \sqrt{2}\sin(\sqrt{2}\Theta_0)\right)^{1/2} \ , & h(\zeta) = 0 \\ u_+(\zeta, m) = \left(\cos\frac{m\zeta}{2} + \frac{1}{m-1}\sin\frac{m\zeta}{2}\right)^{\frac{1}{m}}\left(1 \pm \sqrt{2}\sin(\sqrt{2}\Theta_+)\right)^{1/2} \ , & h(\zeta) = \lambda^2 \ , \end{cases} \tag{48}$$

where

$$\begin{cases} \Theta_- = \frac{(m-1)e^{m\zeta}+1}{\left(e^{\frac{m\zeta}{2}}+\frac{e^{-\frac{m\zeta}{2}}}{m-1}\right)^{\frac{2}{m}}} {}_2F_1\left(1, \frac{m-1}{m}; \frac{m+1}{m}; -\left(\sqrt{m-1}e^{\frac{m\zeta}{2}}\right)^2\right) \\ \Theta_0 = \zeta\, {}_2F_1\left(\frac{1}{m}, \frac{2}{m}; \frac{m+1}{m}; -\frac{1}{m-1}\zeta^m\right) \\ \Theta_+ = \frac{2\sin^{\frac{2}{m}}\left(\frac{m\zeta}{2}+\arctan{(m-1)}\right)\cos\left(\frac{m\zeta}{2}+\arctan{(m-1)}\right)}{m\left(\cos\frac{m\zeta}{2}+\frac{1}{m-1}\sin\frac{m\zeta}{2}\right)^{\frac{2}{m}}} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} + \frac{1}{m}; \frac{3}{2}; \cos^2\left(\frac{m\zeta}{2} + \arctan(m-1)\right)\right) \ . \end{cases} \tag{49}$$

Note that for the case $m = 2$ which is the standard EP case the solution simplifies to

$$\begin{cases} u_-(\zeta, 2) = \sqrt{\left(e^{\zeta} + e^{-\zeta}\right)\left(1 + \sqrt{2}\sin\left(\sqrt{2}\arctan e^{\zeta}\right)\right)} \ , \\ u_0(\zeta, 2) = \sqrt{\left(\zeta + \zeta^{-1}\right)\left(\zeta + \sqrt{2}\zeta\sin\left(\sqrt{2}\mathrm{arctan}\zeta\right)\right)} \ , \\ u_+(\zeta, 2) = \sqrt{(\cos\zeta + \sin\zeta)\left(1 + \sqrt{2}\sin\left(\mathrm{arctanh}\frac{\cos\zeta - \sin\zeta}{\sqrt{2}}\right)\right)} \ . \end{cases} \tag{50}$$

Plots of the solutions u for different values of m and all the other parameters set to the value of unity are given in Figs. (1)-(6). Solutions u_- are real, while u_+ are periodically pure real and pure imaginary. As for u_0, for $m = 2$, it is pure imaginary on the negative half-axis and pure real on the positive half-axis, while for $m > 2$ the solutions display branch cuts. Notice also the diminishing of the amplitudes with increased order of the negative-power nonlinearity.

VII. CONCLUSION

Ermakov-Pinney equations, eiher with standard $m = 2$ or higher-order $m > 2$ nonlinearities, have been treated as dissipative second-order ODEs with the dissipation induced by means of an Abel integrability criterion as discussed in [7]. Using the dynamic invariant of an Ermakov system built with equations of this type, the general solution of these equations is given in the general case of odd Ermakov nonlinearities.

---

[1] P.G. Kevrekidis and Y. Drossinos, Math. Comp. Sim. **74**, 196 (2007).

[2] P.G.L. Leach and K. Andriopoulos, Appl. Anal. Discr. Math. **2**, 146 (2008).

[3] V.P. Ermakov, Univ. Izv. Kiev Ser. III **9**, 1 (1880) and Appl. Anal. Discr. Math. **2**, 123 (2008), translation by A.O. Harin.

[4] J.R. Ray, J.L. Reid, Phys. Lett. A **71**, 317 (1979). J.L. Reid, J.R. Ray, Lett. Math. Phys. **4**, 235 (1980).

[5] W. Sarlet, Phys. Lett. A **82**, 161 (1981).

[6] J.F. Cariñena, J. de Lucas, and M.F. Rañada, Nonlinear superpositions and Ermakov systems, in: F. Cantrijn, M. Crampin, B. Langerock (Eds.), Differential Geometric Methods in Mechanics and Field Theory, Academic Press, 2007. J.F. Cariñena and J. de Lucas, Phys. Lett. A **372**, 5385 (2008).

[7] S.C. Mancas and H.C. Rosu, Integrable dissipative nonlinear second order differential equations via factorizations and Abel equations, arXiv:1212.3636.

[8] H.C. Rosu and O. Cornejo-Pérez, Phys. Rev. E **71**, 046607 (2005). O. Cornejo-Pérez and H.C. Rosu, Prog. Theor. Phys. **114**, 533 (2005). O. Cornejo-Pérez, J. Negro, L.M. Nieto, and H.C. Rosu, Found. Phys. **36**, 1587 (2006).

[9] E. Pinney, Proc. Am. Math. Soc. **1**, 681 (1950).

[10] H. Qin and R.C. Davidson, Phys. Rev. ST Accel. Beams **9**, 054001 (2006).

[11] J.L. Reid, Proc. Am. Math. Soc. **27**, 601 (1971).



FIG. 1: (Color online) $u_{-}(\zeta)$ for the case $m = 2$.



FIG. 2: (Color online) $u_{-}(\zeta)$ for the case $m = 3$ (blue) and $m = 5$ (red).



FIG. 3: (Color online) The modulus square of $u_0(\zeta)$ for the case $m = 2$.



FIG. 4: (Color online) The modulus square of $u_+(\zeta)$ for the case $m = 2$.



FIG. 5: (Color online) The modulus square of $u_+(\zeta)$ for the case $m = 3$.



FIG. 6: (Color online) The modulus square of $u_{+}(\zeta)$ for the case $m = 5$.